

Searles®

MANCOZEB PLUS

Fungicide

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Searles®

MANCOZEB PLUS

Fungicide



ACTIVE CONSTITUENTS:
549g/Kg SULPHUR (S) present as wettable sulphur
235g/Kg MANCOZEB

For control of various Fungal Diseases on some Fruit and Vegetables, Ornamentals and Lawns in the home garden.

The inner bag **MUST NOT** be sold separately.
DO NOT destroy outer box while product still remains.

Net Contents: 200g

Searles®

MANCOZEB PLUS

Fungicide

HOW TO PREPARE:

Product can be applied through most garden sprayers. Add about half the required volume of water to the sprayer and then add the product. Mix or agitate and then add the remaining water. Regular agitation during spraying is necessary to avoid product settling out. Do not store mixed product. Only mix what you can use immediately. Thoroughly cover plants with spray ensuring all leaves are wetted. Avoid excessive runoff.

CAUTION:

Apricots may be damaged due to zinc phytotoxicity. **DO NOT** allow chemical containers or sprays to get into drains, sewers, streams or ponds. **DO NOT** feed grass clippings from treated areas to poultry or animals.

STORAGE AND DISPOSAL:

Store in the closed, original container in a cool, dry place out of the reach of children. Do not store in direct sunlight. Dispose of empty container by wrapping in paper, placing in plastic bag and putting in garbage.

SAFETY DIRECTIONS:

May irritate the eyes, nose, throat and skin. Avoid contact with eyes and skin. **DO NOT** inhale dust or spray mist. Wash hands after use.

Searles®

MANCOZEB PLUS

Fungicide

FIRST AID:

If poisoning occurs, contact a Doctor or Poisons Information Centre (Phone 131 126). Avoid giving alcohol.

J.C. & A.T. Searle Pty. Ltd.
4914 D'Aguilar Highway,
KILCOY. QLD. 4515
www.searles.com.au

NRA Approval No. 55443/1202

BARCODE
9315138007618

HOW TO USE

• **DO NOT** apply if rain is expected within 24 hours or if temperature exceeds 27°C.

Situation	Disease	Rate	How to Apply
AZALEAS, CAMELLIAS	Petal Blight	Mix 50g per 10L of water	Spray when risk of disease is high and repeat spray every 10 days. It is important to thoroughly cover all plant surfaces.
BEANS	Anthrachnose, Leaf Spot, Rust		
BRASSICAS (e.g.: Cabbage, Cauliflower, Broccoli)	Downy Mildew		
CARNATIONS, HOLLYHOCKS, PELARGONIUM, SNAPDRAGONS	Anthrachnose, Leaf Spot, Rust		
CELERY	Leaf Blight		
CUCURBITS (e.g.: Pumpkins, Melons, Cucumber)	Powdery Mildew		
DAHLIAS, GLADIOLI, ZINNIA	Leaf Blight		
GRAPE VINES	Powdery Mildew		
LAWNS	Brown Patch, Dollar Spot (VIC, NSW, SA & WA only)		
LETTUCE	Downy Mildew		
ONION			
ROSES	Black Spot, Powdery Mildew		
POME FRUIT (Apples & Pears)	Black Spot, Powdery Mildew, Bitter Rot		
SEEDLINGS (general)	Damping Off		
STONE FRUIT	Brown Rot, Rust, Shothole, Freckle	Apply at full blossom, petal fall and then 3 week intervals till 2 weeks before harvest.	
TOMATO	Anthrachnose, Early Blight	Spray when risk of disease is high and repeat spray every 10 days. It is important to thoroughly cover all plant surfaces.	

**NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED.
DO NOT PICK POME AND STONE FRUIT, TOMATOES, LETTUCE, GRAPES FOR 14 DAYS AFTER SPRAYING.
DO NOT PICK OTHER EDIBLE CROPS FOR 7 DAYS AFTER SPRAYING.**

Batch No.:

Expiry Date: